

# (Still) Testing for fixed effects in general linear models

Julia Volaufova

*Louisiana State University Health Sciences Center, New Orleans, USA*

## Abstract

Seemingly, testing for fixed effects in linear models with variance-covariance components has been solved for decades. However, even in simple situations such as in fixed one-way model with heteroscedastic variances (a multiple means case of the Behrens-Fisher problem) the questions of statistical properties of various approximations of test statistics are still alive. Here we present a brief overview of approaches suggested in the literature as well as those available in statistical software, accompanied by a simulation study in which the accuracy of p-values is studied.

## Keywords

Mixed linear model, Fixed effects, Heterogeneous variance

## References

- Hrong-Tai Fai, A. and P.L. Cornelius (1996). Approximate F-test of multiple degree of freedom hypotheses in generalized least squares analyses of unbalanced split-plot experiments. *J. Statist. Comput. Simul.* 54, 363–378.
- Giesbrecht, F.G. and J.C. Burns (1985). Two-stage analysis based on mixed model: large-sample asymptotic theory and small-sample simulation results. *Biometrics* 41, 477–486.
- Kenward, G.M. and J.H. Roger (1997). Small sample inference for fixed effects from restricted maximum likelihood. *Biometrics* 53, 983–997.
- Krishnamoorthy, K., F. Lu, and T. Mathew (2007). A parametric bootstrap approach for ANOVA with unequal variances: fixed and random models. *Comput. Statist. Data Anal.* 51, 5731–5742.
- Satterthwaite, F.E. (1946). An approximate distribution of estimates of variance components. *Biometrics Bulletin* 2, 110–114.
- Welch, B.L. (1951). On the comparison of several mean values: an alternative approach. *Biometrika* 38, 330–336.